

Elliptic curves with 3-adic Galois representation surjective mod 3 but not mod 9

Noam D. Elkies¹

Abstract. Let E be an elliptic curve over $\mathbf{Q}$, and $\rho_l : \text{Gal}(\bar{\mathbf{Q}}/\mathbf{Q}) \rightarrow \text{GL}_2(\mathbf{Z}_l)$ its l -adic Galois representation. Serre observed that for $l \geq 5$ there is no proper closed subgroup of $\text{SL}_2(\mathbf{Z}_l)$ that maps surjectively onto $\text{SL}_2(\mathbf{Z}/l\mathbf{Z})$, and concluded that if ρ_l is surjective mod l then it is surjective onto $\text{GL}_2(\mathbf{Z}_l)$. We show that this no longer holds for $l = 3$ by describing a modular curve $\mathcal{X}_9$ of genus 0 parametrizing elliptic curves for which ρ_3 is not surjective mod 9 but generically surjective mod 3. The curve $\mathcal{X}_9$ is defined over $\mathbf{Q}$, and the modular cover $\mathcal{X}_9 \rightarrow X(1)$ has degree 27, so $\mathcal{X}_9$ is rational because 27 is odd. We exhibit an explicit rational function $f \in \mathbf{Q}(x)$ of degree 27 that realizes this cover. We show that for every $x \in \mathbf{P}^1(\mathbf{Q})$, other than the two rational solutions of $f(x) = 0$, the elliptic curves with j -invariant $f(x)$ have ρ_3 surjective mod 3 but not mod 9. We determine all nonzero integral values of $f(x)$, and exhibit several elliptic curves satisfying our condition on ρ_3 , of which the simplest are the curves $Y^2 = X^3 - 27X - 42$ and $Y^2 + Y = X^3 - 135X - 604$ of conductors $1944 = 2^3 3^5$ and $6075 = 3^5 5^2$ respectively.

0. Introduction. Let E be an elliptic curve over $\mathbf{Q}$, and $\rho_l : \text{Gal}(\bar{\mathbf{Q}}/\mathbf{Q}) \rightarrow \text{GL}_2(\mathbf{Z}_l)$ its l -adic Galois representation. Serre observed in 1968 [7, IV, 3.4, Lemma 3] that for $l \geq 5$ there is no proper closed subgroup of $\text{SL}_2(\mathbf{Z}_l)$ that maps surjectively onto $\text{SL}_2(\mathbf{Z}/l\mathbf{Z})$, and concluded that if ρ_l is surjective mod l then it is surjective onto $\text{GL}_2(\mathbf{Z}_l)$. He noted [7, IV, 3.4, Exercise 3] that for $l = 3$ there exists a subgroup $G \subset \text{SL}_2(\mathbf{Z}/9\mathbf{Z})$ such that the restriction to G of the reduction-mod-3 map $\text{SL}_2(\mathbf{Z}/9\mathbf{Z}) \rightarrow \text{SL}_2(\mathbf{Z}/3\mathbf{Z})$ is an isomorphism. The preimage of G in $\text{SL}_2(\mathbf{Z}_3)$ is then a proper closed subgroup that maps surjectively to $\text{SL}_2(\mathbf{Z}/3\mathbf{Z})$. This suggests that there could be curves E for which G is the image of ρ_3 mod 9, making ρ_3 surjective mod 3 but not mod 9. Serre does not raise this question explicitly, and it does not seem to have been addressed elsewhere in the literature; I thank Grigor Grigorov for drawing my attention to it. In this paper we answer the question by showing that there exist infinitely many $j \in \mathbf{Q}$ for which an elliptic curve of j -invariant j must have ρ_3 surjective mod 3 but not mod 9. The simplest examples are $j = 4374$, $j = 419904$, and $j = -44789760$. In general j is the value of a rational function $f(x)$ of degree 27 at all but finitely many $x \in \mathbf{P}^1(\mathbf{Q})$.

Such curves E are parametrized by a modular curve $\mathcal{X}_9 = X(9)/G$. The natural cover $\mathcal{X}_9 \rightarrow X(1)$ has degree 27, and our rational function f arises as the pullback to $\mathcal{X}_9$ of the degree-1 function j on $X(1)$. It is easy to check from the Riemann-Hurwitz formula that $\mathcal{X}_9$ has genus zero. The challenge is to prove that $\mathcal{X}_9$ is defined over $\mathbf{Q}$ and to compute $f(x)$ for some choice of rational coordinate x on $\mathcal{X}_9$. (Once $\mathcal{X}_9$ is known to be defined over $\mathbf{Q}$, it is automatically isomorphic with $\mathbf{P}^1$ over $\mathbf{Q}$, because it supports the rational function j of odd degree.) We prove the rationality in section 1, and compute x using products of Siegel functions in section 2. Such products are modular

¹ Supported in part by NSF grants DMS-0200687 and DMS-0501029.

units on $X(N)$ with known q -expansions; since the cusps of $\mathcal{X}_9$ are not rational we must also find a fractional linear transformation over $\overline{\mathbf{Q}}$ that takes the modular unit to a function x defined over $\mathbf{Q}$. In section 3 we discuss elliptic curves E with j -invariants obtained by specializing f , and use explicit computation of curves $X(9)/H$ with $H \subset G$ to show that all such curves with $x \in \mathbf{P}^1(\mathbf{Q})$ satisfy our condition on ρ_3 except for those with $j = 0$. Finally in section 4 we determine the finite set $f(\mathbf{P}^1(\mathbf{Q})) \cap \mathbf{Z}$, and exhibit some specific elliptic curves E whose j -invariants are these integral values.

1. The group G and the curve $\mathcal{X}_9$. The group $\mathrm{SL}_2(\mathbf{Z}/3\mathbf{Z})$ is generated by

$$S = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad T = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix},$$

the images mod 3 of the standard generators of $\mathrm{SL}_2(\mathbf{Z})$. These generators of $\mathrm{SL}_2(\mathbf{Z}/3\mathbf{Z})$ satisfy

$$S^2 = (ST)^3 = -I, \quad T^3 = I.$$

To lift $\mathrm{SL}_2(\mathbf{Z}/3\mathbf{Z})$ to a subgroup G of $\mathrm{SL}_2(\mathbf{Z}/9\mathbf{Z})$, it is enough to lift S, T to matrices mod 9 satisfying the same relations.² A direct search finds 27 such lifts $\tilde{S}, \tilde{T}$, all equivalent under conjugation in $\mathrm{SL}_2(\mathbf{Z}/9\mathbf{Z})$. These yield 27 choices of lift of $\mathrm{PSL}_2(\mathbf{Z}/3\mathbf{Z})$ to a subgroup $G/\{\pm 1\}$ of $\mathrm{PSL}_2(\mathbf{Z}/9\mathbf{Z}) = \mathrm{Aut}(X(9))$, all conjugate in $\mathrm{Aut}(X(9))$. Hence the quotients of $X(9)$ by these subgroups are all equivalent under $\mathrm{Aut}(X(9))$. We choose

$$\tilde{S} = \begin{pmatrix} 0 & 2 \\ 4 & 0 \end{pmatrix}, \quad \tilde{T} = \begin{pmatrix} 4 & 1 \\ -3 & 4 \end{pmatrix},$$

and set $G = \langle \tilde{S}, \tilde{T} \rangle \subset \mathrm{SL}_2(\mathbf{Z}/9\mathbf{Z})$ and $\mathcal{X}_9 = X(9)/(G/\{\pm 1\})$.

We can then calculate the genus of $\mathcal{X}_9$ by applying the Riemann-Hurwitz formula either to the quotient map $X(9) \rightarrow \mathcal{X}_9$, using the fact that $X(9)$ has genus 10, or to the covering map $\mathcal{X}_9 \rightarrow X(1)$. In the quotient map, each of the three involutions in G has six fixed points, and each of the four 3-element subgroups has three; thus the ramification divisor has degree $3 \cdot 6 + 4 \cdot 6 = 42 = 2|G| + 2(10 - 1)$, whence $\mathcal{X}_9$ has genus 0. The covering map has degree 27 and is unramified except above the cusp $j = \infty$ and the elliptic points $j = 0, j = 1728$. We find that these points have preimages with multiplicities $9^3, 3^8 1^3$, and $12^2 1^3$ respectively (using m^c as a standard shorthand for c preimages of multiplicity m). Hence the ramification divisor has degree $3 \cdot 8 + 8 \cdot 2 + 12 = 52 = 2(27 - 1)$, so again we conclude that $\mathcal{X}_9$ has genus 0.

In particular, each of the cusps of $X(9)$ has trivial stabilizer in G . To explain this, note that the stabilizer in $\mathrm{PSL}_2(\mathbf{Z}/9\mathbf{Z})$ of each cusp of $X(9)$ is conjugate to the group of matrices $\{\pm \begin{pmatrix} 1 & a \\ 0 & 1 \end{pmatrix}\}$; if G had a nontrivial intersection with this group then G would contain $\{\pm \begin{pmatrix} 1 & 3 \\ 0 & 1 \end{pmatrix}\}$, contradicting the requirement that the reduction map $G \rightarrow \mathrm{PSL}_2(\mathbf{Z}/3\mathbf{Z})$ be bijective. It follows that $\mathcal{X}_9$ has $27/9 = 3$ cusps.

² Warning: This approach to lifting $\mathrm{SL}_2(\mathbf{Z}/n\mathbf{Z})$ works only for $n \leq 5$ (including $n = 4$), where $\langle s, t \mid s^2 = (st)^3 = t^n = 1 \rangle$ is a presentation of $\mathrm{PSL}_2(\mathbf{Z}/n\mathbf{Z})$. For $n > 5$ more relations must be checked.

To obtain a model of $\mathcal{X}_9$ defined over $\mathbf{Q}$, we must extend G to a group $G' \subset \mathrm{GL}_2(\mathbf{Z}/9\mathbf{Z})$ such that $G' \triangleright G$ and the determinant map $G'/G \rightarrow (\mathbf{Z}/9\mathbf{Z})^*$ is an isomorphism. For the preimages of the squares in $(\mathbf{Z}/9\mathbf{Z})^*$ we use the invertible multiples of the identity (recall that $-I$ is already in G). It remains to lift $\pm \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}$ from $\mathrm{GL}_2(\mathbf{Z}/3\mathbf{Z})$ to $\mathrm{GL}_2(\mathbf{Z}/9\mathbf{Z})$, and we calculate that the unique choice that normalizes G is $\pm \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}$. Together with the invertible scalar matrices this yields a unique extension G' satisfying our conditions, and thus a unique structure for $\mathcal{X}_9$ as a modular curve over $\mathbf{Q}$.

In general, even if a curve of genus 0 is known to be defined over $\mathbf{Q}$ it need not be isomorphic with $\mathbf{P}^1(\mathbf{Q})$. But in our case the curve supports $\mathbf{Q}$ -rational divisors of odd degree (such as the preimage of the cusp, or indeed the preimage of any rational point on $X(1)$), so $\mathcal{X}_9$ must be isomorphic with $\mathbf{P}^1(\mathbf{Q})$. In the next section we choose an isomorphism, and compute j as a rational function of a rational coordinate on $\mathcal{X}_9$.

While $\mathcal{X}_9$ is defined over $\mathbf{Q}$, its three cusps are not rational. More precisely, they are conjugate over the cyclic cubic extension $K := \mathbf{Q}(\zeta + \zeta^{-1})$, where $\zeta = e^{2\pi i/9}$, a primitive 9th root of unity. To see this, we check that G' contains an element $\pm \begin{pmatrix} a & b \\ 0 & 1 \end{pmatrix}$ for $a = \pm 1$ (when $b = 0$) but no other $a \in (\mathbf{Z}/9\mathbf{Z})^*$. (Since G' contains $\begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}$, it is enough to check this for $a = 4$ and $a = 7$, and then we can multiply by the scalar matrix $a^{-1} \in G'$ to reduce to the corresponding statement for G .) Thus the subgroup of $\mathrm{Gal}(\mathbf{Q}(\zeta)/\mathbf{Q})$ that fixes the cusp $i\infty$ consists of the images of ± 1 under the standard identification of $(\mathbf{Z}/9\mathbf{Z})^*$ with $\mathrm{Gal}(\mathbf{Q}(\zeta)/\mathbf{Q})$. Hence that cusp is defined over K but not over $\mathbf{Q}$, and its conjugates under $\mathrm{Gal}(K/\mathbf{Q})$ must be the other two cusps of $\mathcal{X}_9$.

The fact that the cusps of $\mathcal{X}_9$ are not rational will make our computation of rational functions on $\mathcal{X}_9$ somewhat trickier: we can still expand these functions in powers of q , but the coefficients will in general be contained only in K , not in $\mathbf{Q}$, even for a function in $\mathbf{Q}(\mathcal{X}_9)$.

2. Computing x and $f(x)$. For τ in the upper half-plane $\mathcal{H}$, let $q = e^{2\pi i\tau}$ as usual, and let $q_9 = e^{2\pi i\tau/9}$, a local parameter for $X(9)$ and $\mathcal{X}_9$ at the cusp $\tau = i\infty$, with $q = q_9^9$. The curve $X(3)$ has genus zero with rational coordinate

$$H_3 = \left(\frac{\eta(\tau/3)}{\eta(3\tau)} \right)^3 + 3 = q_9^{-3}(1 + 5q - 7q^2 + 3q^3 + 15q^4 - 32q^5 + \dots);$$

the curve $X(9)$ is a $(\mathbf{Z}/3\mathbf{Z})^3$ cover of $X(3)$ whose function field is obtained from $\mathbf{C}(H_3)$ by adjoining cube roots of $H - 3$, $\zeta^3 H - 3$, and $\zeta^{-3} H - 3$. The group $\mathrm{PSL}_2(\mathbf{Z}/3\mathbf{Z})$ acts on $X(3)$ by the fractional linear transformations of H that permute $\{\infty, 3, 3\zeta^3, 3\zeta^{-3}\}$, which are the transformations that preserve $j = H^3(H^3 + 216)/(H^3 - 27)^3$. (Recall that we set $\zeta = e^{2\pi i/9}$, so 1 and $\zeta^{\pm 3}$ are the cube roots of unity.) To compute $\mathcal{X}_9$, we may lift these transformations to elements of G in the group $\mathrm{PSL}_2(\mathbf{Z}/9\mathbf{Z})$ of automorphisms of $X(9)$ and look for a rational function x of degree 1 on the quotient $\mathcal{X}_9 = X(9)/G$. We could try to find x using our above description of the function field of $X(9)$ as a $(\mathbf{Z}/3\mathbf{Z})^3$ extension of $\mathbf{C}(H_3)$, but this seemed like an unpleasant project. Instead we use modular units on $X(9)$, that is, modular functions whose divisors are supported on the cusps.

For integers a, b we have a function $s_{a,b}$ on $\mathcal{H}$ defined by

$$s(a, b) = q_9^\alpha (1 - \zeta^b q_9^a) \prod_{n=1}^{\infty} (1 - \zeta^b q_9^{9n+a}) (1 - \zeta^{-b} q_9^{9n-a}),$$

where $\alpha = (a^2/9 + a - 9/6)/2 = 18B_2(a/9)$. This is a modular unit, namely the ‘‘Siegel function’’ of [4, p.29] with parameters $(a/9, b/9)$, multiplied by a power of $-\zeta$ that is irrelevant for our purpose. Taking $N = 9$ in [4, p.68, Thm. 4.1], we see that a product $F = \prod_r s(a_r, b_r)^{m_r}$ of such functions is a modular function on $X(9)$ if and only if the (a_r, b_r) and m_r satisfy the ‘‘quadratic relations’’

$$\sum_r m_r a_r^2 \equiv \sum_r m_r b_r^2 \equiv \sum_r m_r a_r b_r \equiv 0 \pmod{9}.$$

Note that this condition depends only on the $(a_r, b_r) \pmod{9}$, and is invariant under changing (a_r, b_r) to $(-a_r, -b_r)$ for some r ; Indeed $s(-a, -b)$ is a scalar multiple of $s(a, b)$, as is $s(a', b')$ if $(a', b') \equiv (a, b) \pmod{9}$.

The pairs $\{(a, b), (-a, -b)\} \in (\mathbf{Z}/9\mathbf{Z})^2$ with at least one of a, b not divisible by 3 correspond bijectively with the cusps of $X(9)$: the cusps are orbits of $\Gamma(9)$ acting on $\mathbf{P}^1(\mathbf{Q})$, and if $\gcd(a, b) = \gcd(a', b') = 1$ for some integers a, b, a', b' then a/b and a'/b' are in the same orbit if and only if $(a', b') \equiv (a, b)$ or $(-a, -b) \pmod{9}$. This labeling is consistent with the action of $\mathrm{PSL}_2(\mathbf{Z}/9\mathbf{Z})$ on cusps, and dual to its action on the modular units $s(a, b)$ (up to μ_{18} factors). We choose one of the three orbits of the action of $G/\{\pm 1\}$ on the cusps of $X(9)$, and take (a_r, b_r) corresponding to the twelve cusps in the orbit. We check that these satisfy the quadratic relations with $m_r = 1$, so the product F of the twelve functions $S(a_r, b_r)$ is a modular function on $X(9)$ whose divisor is invariant under G .

There is then a homomorphism $\chi : G \rightarrow \mathbf{C}^*$ such that $F(g(\tau)) = \chi(g)F(\tau)$ for all $g \in G$ and $\tau \in \mathcal{H}$. We claim that χ is trivial. Indeed, if it were nontrivial we could take for g a 3-cycle in G , and for τ a preimage of a fixed point on $X(9)$ of g , and conclude that τ is a zero or pole of F , which is impossible because τ is not a cusp. Therefore χ is trivial as claimed, whence F descends to a function on $\mathcal{X}_9$ whose only poles or zeros are at the cusps.

We find that F has a simple pole at one cusp, a simple zero at another, and neither zero nor pole at the third. In particular, F is a function of degree 1 on $\mathcal{X}_9$. Choosing the orbit of $(a, b) = (1, 0)$, corresponding to the cusp $i\infty$, we find that F has a pole at that cusp, and calculate the q -expansion

$$F = q_9^{-1} - 1 + c_4 q_9 + c_1 q_9^2 + (c_2 + 2) q_9^3 + c_4 q_9^4 \cdots,$$

where $c_m := \zeta^m + \zeta^{-m}$, a unit in K . The q -expansions of the products corresponding to the other two orbits then let us recognize those products as fractional linear transformations of F , namely $1/(F - c_2 + 1)$ and $(1 - c_2)/F$.

It follows that F takes the values 0 and $1 - c_2$ on the other two cusps of $\mathcal{X}_9$. Therefore j , considered as a rational function of F , must have its poles at $F = 0, 1 - c_2, \infty$,

each with multiplicity 9. Indeed we compute that $F^9(F - c_2 + 1)^9 j$ is a polynomial of degree 27 in F to the accuracy allowed by our q -expansions (which extend far enough beyond the constant term of that polynomial to provide a sanity check on our computations).

It remains to find a fractional linear transformation with coefficients in K that, when applied to F , yields a coordinate x on $\mathcal{X}_9$ such that $j \in \mathbf{Q}(x)$. Thus x must map the three cusps to a $\text{Gal}(K/\mathbf{Q})$ orbit in K . We may choose any ordered orbit, and then x is determined uniquely, because PGL_2 acts simply 3-transitively on $\mathbf{P}^1$. (The order must be consistent with the Galois action on the cusps.) We then apply a $\text{PGL}_2(\mathbf{Q})$ transformation so that the map $\mathcal{X}_9 \rightarrow X(1)$ is represented by a rational function with small coefficients. This leads us to

$$\begin{aligned} x &= \frac{-c_1 F + 1 - c_2}{F - c_1 + 3(1 - c_2)} \\ &= -c_1 + (2c_2 + c_4)q_9 + 3(1 - c_1)q_9^2 + (6 - 7c_1 + c_2)q_9^3 + (15 - 16c_1 + 7c_2)q_9^4 \cdots, \end{aligned}$$

when $j = f(x)$ with

$$\begin{aligned} f(x) &= -\frac{3^7(x^2 - 1)^3(x^6 + 3x^5 + 6x^4 + x^3 - 3x^2 + 12x + 16)^3(2x^3 + 3x^2 - 3x - 5)}{(x^3 - 3x - 1)^9} \\ &= 1728 - \frac{3^3 A^2(x) B^2(x) (2x^3 - 3x^2 + 4)}{(x^3 - 3x - 1)^9}, \end{aligned}$$

where $A(x)$, $B(x)$ are the sextic polynomials

$$\begin{aligned} A(x) &= x^6 + 6x^5 + 4x^3 + 12x^2 - 18x - 23, \\ B(x) &= 7x^6 + 24x^5 + 18x^4 - 26x^3 - 33x^2 + 18x + 28. \end{aligned}$$

3. The elliptic curves parametrized by $\mathcal{X}_9$. Now let $E/\mathbf{Q}$ be an elliptic curve with j -invariant $f(x)$ for some $x \in \mathbf{P}^1(\mathbf{Q})$. Assume that its 3-adic Galois representation ρ_3 is surjective mod 3. Then its image mod 9 is (a conjugate of) the proper subgroup G' of $\text{GL}_2(\mathbf{Z}/9\mathbf{Z})$, because this image is contained in G and its determinant maps surjectively to $\mathbf{Z}_3^*$. In particular, ρ_3 is surjective mod 3 but not mod 9, as desired.

We claim that the mod-3 condition on ρ_3 is satisfied by except at $x = \pm 1$, the points at which $f(x) = 0$, the j -invariant of an elliptic curve with complex multiplication.

For any elliptic curve $E/\mathbf{Q}$, the representation ρ_3 is surjective mod 3 if and only if the intersection of its image in $\text{PGL}_2(\mathbf{Z}/3\mathbf{Z})$ with $\text{PSL}_2(\mathbf{Z}/3\mathbf{Z})$ is contained in the 4-element normal subgroup or in a cyclic subgroup of order 3. In the former case, the j -invariant j_E of E is a cube, and this necessary condition is also sufficient unless $j_E = 0$. If $f(x) \neq 0$ then $f(x)$ is a cube if and only if $3(2x^3 + 3x^2 - 3x - 5) = z^3$ for some $z \in \mathbf{Q}$. This curve of genus 1 has no rational points due to a 3-adic obstruction: we have

$$2x^3 + 3x^2 - 3x - 5 = 3(x + 1)^3 - (x + 2)^3,$$

so the 3-adic valuation of $3(2x^3 + 3x^2 - 3x - 5)$ is never divisible by 3. Thus $f(x)$ is never a nonzero cube. The latter case holds if and only if E admits a rational

3-isogeny. Such x are parametrized by a curve $\mathcal{X}'_9 = \mathcal{X}_9/\langle \tilde{T} \rangle$, the quotient of $X(9)$ by a 3-cycle in G . This curve has genus 3, so there are only finitely many such x by Mordell-Faltings. For a general curve of genus 3 it is not known how to provably list all the rational points. But here we are lucky: we can give a rational map of degree 3 from $\mathcal{X}'_9$ to the elliptic curve $Y^2 + Y = X^3$, which is known to have rank zero. Pulling back each of this curve's three rational points to $\mathcal{X}'_9$ then yields a set of $\bar{\mathbf{Q}}$ -rational points that must contain all the $\mathbf{Q}$ -rational ones. To complete the proof of our claim we shall observe that there are only two $\mathbf{Q}$ -rational points, one for each of $x = \pm 1$.

A simple model for $\mathcal{X}'_9$ is

$$3z_2^3 = z_1(z_1^3 + 3z_1^2 - 6z_1 + 1), \quad (1)$$

and the map to $Y^2 + Y = X^3$ can be given by

$$(X : Y : 1) = (z_1(z_1 + 1)z_2 : 3z_1^2 : z_2^3). \quad (2)$$

The rational points on $\mathcal{X}'_9$ are $(0, 0)$ and the point at infinity. The equation (1) for $\mathcal{X}'_9$ shows that this curve has an automorphism σ of order 3 that fixes z_1 and multiplies z_2 by a cube root of unity. This automorphism arises from the element $3\tilde{T} - 2 = \begin{pmatrix} 1 & 3 \\ 0 & 1 \end{pmatrix}$ of $\mathrm{PSL}_2(\mathbf{Z}/9\mathbf{Z})$, which commutes with T and thus descends from $\mathrm{Aut}(X(9))$ to an automorphism of $\mathcal{X}'_9$. The quotient of $\mathcal{X}'_9$ by $\langle \sigma \rangle$ is the z_1 -line, which covers the j -line $X(1)$ with degree 36 and the curve $X_0(3)$ with degree 9. The latter map can be realized by the rational function $27((z_1^3 + 3z_1^2 - 6z_1 + 1)/(z_1^3 - 6z_1^2 + 3z_1 + 1))^3$.

We next outline the computation of the model (1) and the map (2). We begin with the modular units

$$s(1, 0)s(4, 6)s(4, 3), \quad s(4, 0)s(7, 6)s(7, 3), \quad s(7, 0)s(1, 6)s(1, 3). \quad (3)$$

The first of these corresponds to the orbit of the cusp $i\infty$ under the 3-cycle $\tilde{T}$, and the others are obtained by multiplying $\{(1, 0), (4, 6), (4, 3)\}$ by 4 and 7 mod 9. We calculate that these products do not satisfy the quadratic relations, but the quotient of any two of them does, giving a rational function of degree 3 on $\mathcal{X}'_9$. Using the q -expansions we find that the three functions in (3) are linearly dependent, and thus that their pairwise quotients are all related by fractional linear transformations over K . As we did for x , we find two linear combinations of the functions in (3) whose quotient is a degree-3 function on $\mathcal{X}'_9$ defined over $\mathbf{Q}$. We call this function

$$y = -c_2 + (c_4 - c_2 + 3)q_9 + (3c_4 - 6c_7 + 9)q_9^2 + (10c_4 - 22c_7 + 27)q_9^3 \cdots,$$

and use the q -expansions to find the coefficients of a polynomial identity $P(x, y) = 0$ of bidegree $(3, 4)$ in (x, y) .

This is a model for $\mathcal{X}'_9$ in $\mathbf{P}^1 \times \mathbf{P}^1$, but it is not smooth. We find that it has three double points, at $x = y = -c_1, -c_2, -c_4$. Thus the holomorphic differentials on $\mathcal{X}'_9$ are the forms $Q(x, y) dx/P_y$ for Q in the 3-dimensional space of polynomials of bidegree $(1, 2)$ that vanish at the three singularities. We interpret these as modular cuspforms of

weight 2 on $\mathcal{X}'_9$ by writing $q dx/dq$ in place of dx , and find a basis for the space of such cuspforms:

$$\begin{aligned}\varphi_1 &= c_4q - 3q^3 - 2c_2q^4 - c_1q^7 + 6q^{12} + 5c_2q^{13} + 4c_1q^{16} - 7c_4q^{19} + 3q^{21} \dots, \\ \varphi_2 &= q + (c_4 - c_1)q^2 + q^4 + (2c_2 + c_4)q^5 + 2q^7 + (c_2 + c_4)q^8 - 3q^{10} \dots, \\ \varphi_3 &= (c_4 - c_2)q - 3q^2 + (c_2 - c_1)q^4 + 3q^5 + (2c_1 - 2c_4)q^7 + 3q^8 + (3c_2 - 3c_4)q^{10} \dots.\end{aligned}$$

The affine model (1) is then obtained by taking $(z_1 : z_2 : 1) = (\varphi_2 + \varphi_1 : \varphi_3 : 2\varphi_2 - \varphi_1)$. The map (2) was obtained by integrating the CM form φ_1 ; it can also be seen in (1) by writing the cubic factor as $(z_1 + 1)^3 - 9z_1$.

4. Numerical examples. Besides $f(\pm 1) = 0$, there are seven other integers obtained by evaluating $f(x)$ at points of $\mathbf{P}^1(\mathbf{Q})$ of small height. For each of those, we list x , $f(x)$, and one of the elliptic curves E of j -invariant $f(x)$ and minimal conductor N :

x	$j = f(x)$	E	N
1/0	4374	$[0, 0, 0, -27, -42]$	$2^3 3^5$
-2	419904	$[0, 0, 0, -162, 792]$	$2^8 3^5$
0	-44789760	$[0, 0, 1, -135, -604]$	$3^5 5^2$
-1/2	15786448344	$[0, 0, 0, -5427, 153882]$	$2^5 3^5$
2	24992518538304	$[0, 0, 0, -201042, 34695912]$	$2^8 3^5 17^2$
-3/2	-92515041526500	$[0, 0, 0, -1126035, 459913278]$	$2^3 3^5 19^2$
-1/3	-70043919611288518656	$[0, 0, 1, -1127379978, -14569799990728]$	$3^5 97^2 101^2$

The smallest conductor here is $1944 = 2^3 3^5$, still too large to appear in Cremona's published tables [2] of curves of conductor ≤ 1000 . But Cremona has pursued his computations up to 10^5 and beyond (see [3] for the status as of mid-2006), enough to find our first curve as well as those of conductors $6075 = 3^5 5^2$, $7776 = 2^5 3^5$, and $62208 = 2^8 3^5$. Curves with $j = 4374$, $j = 419904$, and $j = 15786448344$ already appeared in the tables of elliptic curves with good reduction away from 2 and 3, compiled in 1966 by F.B. Coghlan and published as "Table 4" in [1, p.123]: see rows 52, 84, and 86 of "Table 4a" [1, p.125], and [1, p.75] for the attribution to Coghlan.

A search up to height 256 found no more integral values of $f(x)$. Since f has three distinct poles, there can be only finitely many x for which $f(x) \in \mathbf{Z}$. We claim that in fact we have found them all. Suppose $x = m/n$ in lowest terms. The resultant of the numerator and denominator of $f(x)$ is 3^{486} , so when we write $f(m/n)$ as the quotient of homogeneous polynomials in m, n the denominator $(m^3 - 3mn^2 - n^3)^9$ must be $\pm 3^A$ for some A . Thus $m^3 - 3mn^2 - n^3 = \pm 1$ or ± 3 , because the only $(m, n) \in \mathbf{Z}^2$ for which $9 \mid m^3 - 3mn^2 - n^3$ are those for which $3 \mid m$ and $3 \mid n$. The only cases of $m^3 - 3mn^2 - n^3 = \pm 3$ are $x = 1, -2, -1/2$, because these are the only rational points on the elliptic curve $x^3 - 3x - 1 = 3z^3$ (isogenous with the cubic Fermat curve). For $m^3 - 3mn^2 - n^3 = \pm 1$ one must work harder because the elliptic curve $x^3 - 3x - 1 = z^3$ has rank 1. Fortunately this work was already done by Ljunggren, who proved in 1942 that the Thue equation $m^3 - 3mn^2 - n^3 = 1$ has only the six solutions $(1, 0)$, $(0, -1)$, $(-1, 1)$, $(2, 1)$, $(1, -3)$, $(-3, 2)$, corresponding to values of x already listed above. (See [5, §2], cited by Nagell [6] who also notes the connection between $X^3 + Y^3 = Z^3$

and $m^3 - 3mn^2 - n^3 = 3$.) The solutions of $m^3 - 3mn^2 - n^3 = -1$ are obtained from these six by changing each (m, n) to $(-m, -n)$, which yields the same values of $x = m/n$. Thus our list contains all the nonzero integral j -invariants of elliptic curves parametrized by $\mathcal{X}_9$.

References

- [1] Birch, B.J., Kuyk, W., ed.: *Modular Functions of One Variable IV*. Lect. Notes in Math. **476**, 1975.
- [2] Cremona, J.E.: *Algorithms for modular elliptic curves*. Cambridge University Press, 1992.
- [3] Cremona, J.E.: The elliptic curve database for conductors to 130000, Lecture Notes in Computer Science 4076 (Proceedings of ANTS-7, 2006; F.Hess, S.Pauli, and M.Pohst, ed.), 11–29.
- [4] Kubert, D.S., Lang, S.: *Modular Units*. New York: Springer 1991.
- [5] Ljunggren, W.: Einige Bemerkungen über die Darstellung ganzer Zahlen durch binäre kubische Formen mit positiver Diskriminante, *Acta math.* **75** (1942), 1–21.
- [6] Nagell, T.: Sur un type particulier d’unités algébriques, *Arkiv för Math.* **8** (1969), 163–184.
- [7] Serre, J.-P.: *Abelian l -adic Representations and Elliptic Curves*. Wellesley, Mass.: A.K. Peters, 1997.